ALL DAY ALLERGY-D- cetirizine hcl, pseudoephedrine hcl tablet, extended release L. Perrigo Company

Perrigo All Day Allergy-D Drug Facts

Active ingredients (in each extended release tablet)

Cetirizine HCl 5 mg

Pseudoephedrine HCl 120 mg

Purpose

Antihistamine

Nasal Decongestant

Uses

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- thyroid disease
- diabetes
- glaucoma
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are

taking tranquilizers or sedatives.

When using this product

- do not use more than directed
- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- an allergic reaction to this product occurs. Seek medical help right away.
- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

If pregnant or breast-feeding:

- if breast-feeding: not recommended
- if pregnant: ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not break or chew tablet; swallow tablet whole

adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

store between 20° to 25°C (68° to 77°F)

Inactive ingredients

colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredients in Zyrtec-D®

Original Prescription Strength

All Day Allergy-D

Cetirizine Hydrochloride and Pseudoephedrine Hydrochloride Extended Release Tablets, 5 mg/120 mg

Antihistamine/Nasal Decongestant

Allergy & Congestion

INDOOR & OUTDOOR ALLERGIES

12 hour

12 HOUR RELIEF OF:

Sneezing

Itchy, Watery Eyes

Runny Nose


Itchy Throat or Nose

Sinus Pressure

Nasal Congestion

Actual Size

12 Extended Release Tablets



Drug Facts

Active ingredients

Purpose (in each extended release tablet)

Cetirizine HCI 5 mg......Antihistamine Pseudoephedrine HCI 120 mg.....Nasal Decongestant .Antihistamine

Uses

■ temporarily relieves these symptoms due to hay

Drug Facts (continued)

fever or other upper respiratory allergies:

- runny nose ■ itchy, watery eyes ■ sneezing ■ itching of the nose or throat
 ■ nasal congestion
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through



17653 55 0

Drug Facts (continued)

Warnings

Do not use

- if you have ever had an allergic reaction to this product or any of its ingredients or to an antihistamine containing hydroxyzine.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doct or or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease thyroid disease diabetes
- glaucoma high blood pressure
- trouble urinating due to an enlarged prostate gland ■ liver or kidney disease. Your doctor should determine if you need a different dose.

Ask a doctor or pharmacist before use if you are taking tranquilizers or sedatives.

When using this product

- do not use more than directed
- drowsiness may occur avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

■ an allergic reaction to this product occurs. Seek medical help right away.

Drug Facts (continued)

- you get nervous, dizzy, or sleepless
- symptoms do not improve within 7 days or are accompanied by fever

If pregnant or breast-feeding:

- ■if breast-feeding: not recommended
- ■if pregnant: ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not break or chew tablet; swallow tablet whole

adults and children 12 years and over	take 1 tablet every 12 hours; do not take more than 2 tablets in 24 hours.
adults 65 years and over	ask a doctor
children under 12 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

store between 20° to 25°C (68° to 77°F)

Inactive ingredients

colloidal silicon dioxide, hydroxypropyl cellulose, hypromellose, lactose monohydrate, magnesium stearate, microcrystalline cellulose, talc, yellow iron oxide





Drug Facts (continued)

(Questions or comments? 1-800-719-9260

Distributed By **Perrigo**® Allegan, MI 49010 perrigo.com



ALL DAY ALLERGY-D

cetirizine hcl, pseudoephedrine hcl tablet, extended release

Product Information Product Type HUMAN OTC DRUG Item Code (Source) NDC:0113-2176 Route of Administration ORAL

Active Ingredient/Active Moiety Basis of Strength Ingredient Name Strength CETIRIZINE HYDRO CHLO RIDE (UNII: 640047KTOA) (CETIRIZINE -CETIRIZINE HYDROCHLORIDE 5 mg UNII:YO7261ME24) PSEUDO EPHEDRINE HYDRO CHLO RIDE (UNII: 6 V9 V2RYJ8 N) (PSEUDO EPHEDRINE **PSEUDOEPHEDRINE** 120 mg - UNII:7CUC9DDI9F) HYDROCHLORIDE

Inactive Ingredients		
Ingredient Name	Strength	
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)		
HYPROMELLOSES (UNII: 3NXW29 V3WO)		
LACTO SE MO NO HYDRATE (UNII: EWQ57Q8 I5X)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
CELLULO SE, MICRO CRYSTALLINE (UNII: OP1R32D61U)		
TALC (UNII: 7SEV7J4R1U)		
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)		

Product Characteristics			
Color	WHITE (one side white one side light yellow)	Score	no score
Shape	ROUND	Size	12mm
Flavor		Imprint Code	5029;5;120
Contains			

Packaging				
# Item C	Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:0113-	2176-62	4 in 1 CARTON	09/06/2011	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2 NDC:0113-	2176-53	2 in 1 CARTON	09/26/2013	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077170	09/06/2011	

Labeler - L. Perrigo Company (006013346)

Revised: 12/2018 L. Perrigo Company